arXiv:1301.4740v2 [cond-mat.str-el] 23 Jan 2013

Frustration effect on magnetic excitations in a two-leg spin-ladder system

Takanori Sugimoto,[1, 2, *] Michiyasu Mori,[1, 2] Takami Tohyama,[3] and Sadamichi Maekawa[1, 2]
[1] *Advanced Science Research Center, Japan Atomic Energy Agency, Tokai, Ibaraki, 319-1195, Japan*
[2] *CREST, Japan Science and Technology Agency,*
3-Sanbancho, Chiyoda-ku, Tokyo, 102-0075, Japan
[3] *Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto, 606-8502, Japan*

(Dated: January 24, 2013)

We theoretically study the magnetic excitations in a frustrated two-leg spin-ladder system, in which antiferromagnetic exchange interactions act on the nearest-neighbor and next-nearest-neighbor bonds in the leg direction, and on the nearest-neighbor bonds in the rung direction. A dynamical spin correlation function at zero temperature is calculated by using the dynamical density-matrix renormalization-group method for possible magnetic phases, i.e., columnar-dimer and rung-singlet phases. The columnar-dimer phase is characterized by multi-spinon excitations with spin gap, while the rung-singlet phase is dominated by the triplet excitation in the rung direction. It is found that a major difference between these two phases appears in the spectral weight of magnetic excitations, in particular, of the bonding and anti-bonding modes in the rung direction. Therefore, we can distinguish one phase from the other by the difference of the spectral weight. Furthermore, we examine frustration effect on both modes in the rung-singlet phase with a perturbation theory from the strong coupling limit. It is shown that the anti-bonding mode is stable against the frustration, and a wavenumber with minimum excitation energy is shifted from the commensurate to incommensurate ones. On the other hand, the bonding mode is merged into the continuum excitation of multiple triplet excitations by increasing the frustration. By comparing our results with inelastic neutron scattering experiments for $\mathrm{BiCu_2PO_6}$, the magnitude of the magnetic exchange interactions and the ground state will be determined.

I. INTRODUCTION

Spinon is an elementary excitation characteristic to the strongly correlated electron systems in low dimensions[1], and was observed by angle-resolved photoemission spectroscopy[2] and neutron scattering[3]. It is a manifestation of electron fractionalization called spin-charge separation, in which the electronic excitation is split into the spin degree of freedom (spinon) and the charge one (holon), instead of a quasiparticle. In the frustrated spin systems with exotic ground states such as spin-liquid and valence-bond-solid states, the spinon is associated with a topological excitation similar to the magnetic domain wall[4]. In contrast to the collective excitation in a magnetic long-ranged ordered state, i.e., magnon, the spinon carries a fractional quantum number, spin 1/2, accompanied by a spin string. Because of this topological nature, two spinons make a bound state called triplon in some cases[4,5].

The frustrated two-leg spin-ladder system, i.e., J_1-J_2-J_p model (Fig. 1), provides those exotic magnetic states, since this model bridges two different systems: the frustrated spin chain, i.e., J_1-J_2 model and the unfrustrated spin ladder, i.e., J_1-J_p model. Here, J_1 and J_2 are the nearest and the next-nearest antiferromagnetic exchange interactions, respectively, and J_p is the antiferromagnetic interaction in the rung direction. The ground-state phase diagram has been examined numerically[6], and has two phases; the columnar-dimer (CD) phase, which continuously merges to the dimer phase of the frustrated chain system (J_1-J_2 model) in the zero limit of J_p, and the rung-singlet (RS) phase, which continuously changes to the ground state of the unfrustrated ladder system (J_1-J_p model) in the zero limit of J_2. Note that the phase transition between the RS and the CD phases is of the Ising-type[6]. On the other hand, the excited states of the J_1-J_2-J_p model will be non-trivial, since the frustration leads to the confinement of spinons and triplons[4]. The two limiting cases of the J_1-J_2 and J_1-J_p models are involved in the J_1-J_2-J_p model. Therefore, this frustrated two-leg spin-ladder is useful for understanding the frustration effect on magnetic excitations. Several properties of the two limiting cases are summarized below.

The J_1-J_2 model has two ground states: gapless spin-liquid and gapped dimer phases. Between these phases, Kosterlitz-Thouless transition occurs by changing the ratio of the exchange interactions, J_2/J_1[7–9]. In the spin-liquid phase, spin correlation function exhibits a critical behavior and has no long-range order. The excitations are understood well with the spinon picture, which is obtained by the Jordan-Wigner transformation of the spin operators. In particular, the triplet excitation is dominated by two-spinon excitation[10,11]. In the dimer phase, neighboring two spins constitute a singlet dimer, whose dimer correlation remains finite in the limit of infinite distance[12,13]. The ground state is doubly degenerated because of two choices to form a singlet dimer with either a right neighboring site or a left one.

The intensive studies on the unfrustrated ladder system, i.e., the J_1-J_p model, show that one-dimensional spin-liquid phase is fragile with respect to the interchain interaction[14]. The RS phase appears in the ground state[14], where two spins connecting a rung bond constitute a singlet dimer and there is no degeneracy. Thus, the RS phase is another gapped phase different from the dimer phase. The magnetic excitation is dominated by triplon, which is a bosonic quasiparticle defined as the triplet excitations from the RS dimers[15,16]. It is also interpreted as a bound state of two spinons[3–5]. The triplon always appears as an anti-bonding mode in the rung direction due to the parity changes between the ground state and the triplon excited state[17]. On the other hand, the excitation with even number of triplons appears as a bonding mode in the rung direction. The bound states of two triplons are important for low-energy excitation in the bonding modes, since the intensity is much larger than the two-triplon continuum[16].

In this paper, we investigate the magnetic excitations in the J_1-J_2-J_p model. Our main purpose is to make clear the frustration effects by J_2 on the excitation spectra, which are observed by inelastic neutron scattering for $\mathrm{BiCu_2PO_6}$ (Ref. 18-21). The compound should be compared with the unfrustrated two-leg spin-ladder compound, $\mathrm{SrCu_2O_3}$ (Ref. 22). Not only the dispersion relation but also the spectral weight is crucial for analyzing experimental data. Thus, we calculate dynamical spin correlation function (DSCF) by dynamical density-matrix renormalization-group (dynamical DMRG) method. Furthermore, we propose how to determine the ground-state phase and how to estimate the interaction J_1, J_2 and J_p in $\mathrm{BiCu_2PO_6}$, by combining the DSCF and inelastic-neutron-scattering data. Our another purpose is to clarify difference between two phases, i.e., CD and RS phases, by the DSCF and to obtain relation of the exchange energies and the spectrum.

The rest of the paper is organized as follows. First, we present the model Hamiltonian of the frustrated two-leg spin-ladder system and method to calculate DSCF by dynamical DMRG in Sec. II. We also prepare some notations for a perturbation theory to explain behaviors of the excitations. In Sec. III, we show the numerical results and some interpretations on magnetic excitations. We discuss the frustration effects on the excitation. The ways to estimate the magnitude of the exchange interactions are also discussed. Finally, we summarize the results and discuss consistency with previous studies in Sec. IV.

II. MODEL AND METHOD

The model Hamiltonian of the frustrated two-leg spin-ladder system in Fig. 1 is defined as

$$\mathcal{H} = \sum_j J_p h_j^{(p)} + J_1 h_j^{(1)} + J_2 h_j^{(2)} \tag{1}$$

with

$$h_j^{(p)} = \boldsymbol{S}_{j,\mathrm{u}} \cdot \boldsymbol{S}_{j,\mathrm{l}}, \tag{2}$$

$$h_j^{(1)} = \boldsymbol{S}_{j,\mathrm{u}} \cdot \boldsymbol{S}_{j+1,\mathrm{u}} + \boldsymbol{S}_{j,\mathrm{l}} \cdot \boldsymbol{S}_{j+1,\mathrm{l}}, \tag{3}$$

$$h_j^{(2)} = \boldsymbol{S}_{j,\mathrm{u}} \cdot \boldsymbol{S}_{j+2,\mathrm{u}} + \boldsymbol{S}_{j,\mathrm{l}} \cdot \boldsymbol{S}_{j+2,\mathrm{l}}, \tag{4}$$



FIG. 1. A frustrated two-leg spin-ladder system. Balls represent spin-$\frac{1}{2}$ sites. The nearest- and next-nearest-neighbor interactions in the leg direction, J_1 and J_2, are associated with solid and dashed lines, respectively. Double solid line denotes the nearest-neighbor interaction in the rung direction J_p.

where $J_1(>0)$ and $J_2(>0)$ are the magnitude of antiferromagnetic nearest-neighbor and next-nearest-neighbor exchange interactions, respectively, and $J_p(>0)$ is that of antiferromagnetic nearest-neighbor interaction in the rung direction. $\boldsymbol{S}_{j,\mathrm{u(l)}}$ is the $S=1/2$ spin operator on j site in the upper (lower) chain.

To investigate magnetic excitations in the frustrated ladder system, we calculate the z component of DSCF at zero temperature given by

$$\begin{aligned}\chi(\boldsymbol{q},\omega) &= -\frac{1}{\pi}\Im\int_0^\infty dt e^{i\omega t}\langle 0|S^{z\dagger}(\boldsymbol{q},t)S^z(\boldsymbol{q},0)|0\rangle \\ &= -\lim_{\gamma\to 0}\frac{1}{\pi}\Im\langle 0|S^z(-\boldsymbol{q})\frac{1}{\omega-\mathcal{H}+\epsilon_0+i\gamma}S^z(\boldsymbol{q})|0\rangle,\end{aligned} \tag{5}$$

where $|0\rangle$ is the ground state. The x (y) component equals to the z component because of no anisotropy in the model Hamiltonian (1). To obtain the excitation spectra, we set the infinitesimal value γ to be 0.1, which is enough to examine the qualitative behaviors of the spin excitations in this system. The momentum representation of the spin operator with the open boundary condition is given by,

$$S^z(\boldsymbol{q}) = \sqrt{\frac{2}{N+1}}\sum_j \sin(q_x j) S^z_{j,q_y}, \tag{6}$$

with

$$S^z_{j,q_y=0} = \frac{1}{2}(S^z_{j,\mathrm{u}} + S^z_{j,\mathrm{l}}), \tag{7}$$

$$S^z_{j,q_y=\pi} = \frac{1}{2}(S^z_{j,\mathrm{u}} - S^z_{j,\mathrm{l}}), \tag{8}$$

where x axis is set to the chain direction and y axis is in the rung direction. N is the rung number of the system, namely the total number of spin sites is $2N$.

We use the dynamical DMRG method[23,24] to obtain the DSCF. This method requires three target states: $|0\rangle$, $S^z(\boldsymbol{q})|0\rangle$, and $[\omega-\mathcal{H}+\epsilon_0+i\gamma]^{-1}S^z(\boldsymbol{q})|0\rangle$. The last target state, so-called the correction vector, is obtained by using a modified conjugate gradient method in the dynamical DMRG algorithm.

We check the convergence of the ground-state energy for the DMRG truncation number m. We confirm that the ground-state energy per site converges within a numerical error less than 10^{-3} for $m = 150$ and use this value to obtain magnetic excitation spectra in a 32-rung frustrated ladder system.

A perturbation analysis with resonated valence bond (RVB) picture is also used to obtain the ground state and low-energy excited states. We can rewrite the perturbed Hamiltonian, $h_j^{(R)}$ $(R = 1, 2)$, by the projection operators onto RVB configurations. The RVB picture helps us to handle complicated spin-configurations easier (see Appendix). We prepare the notations for singlet pair, $[j,k]_p$, and triplet one with $S^z = 0$, $\{j,k\}_p$, of two spins on j and k sites $(j < k)$ as follows,

$$[j,k]_{p=\mathrm{i,u,l}} = \frac{1}{\sqrt{2}}(|\uparrow\rangle_{j,a_p}|\downarrow\rangle_{k,b_p} - |\downarrow\rangle_{j,a_p}|\uparrow\rangle_{k,b_p}), \quad (9)$$

$$\{j,k\}_{p=\mathrm{i,u,l}} = \frac{1}{\sqrt{2}}(|\uparrow\rangle_{j,a_p}|\downarrow\rangle_{k,b_p} + |\downarrow\rangle_{j,a_p}|\uparrow\rangle_{k,b_p}), \quad (10)$$

where the index p denotes the position of singlet or triplet bond, i.e., $p = \mathrm{i}$ indicates an inter-chain bond, and $p = \mathrm{u(l)}$ means a bond in the upper (lower) chain. The local spin state on j site in the upper (lower) chain is given by $|\uparrow\rangle_{j,\mathrm{u(l)}}$ or $|\downarrow\rangle_{j,\mathrm{u(l)}}$. Then, $a_\mathrm{i} = \mathrm{u}$ and $b_\mathrm{i} = \mathrm{l}$ for $p = \mathrm{i}$, while $a_\mathrm{u} = b_\mathrm{u} = \mathrm{u}$ $(a_\mathrm{l} = b_\mathrm{l} = \mathrm{l})$ for $p = \mathrm{u(l)}$.

Below, the index p is omitted for the intra-chain, if it is obvious. We define two types of singlet configurations between j and k rungs as,

$$\langle j,k\rangle_\mathrm{r} = [j,j]_\mathrm{i}[k,k]_\mathrm{i}, \quad (11)$$

$$\langle j,k\rangle_\mathrm{l} = [j,k]_\mathrm{u}[j,k]_\mathrm{l}. \quad (12)$$

By using these expression, $S^z = 0$ states in total $S = 0$ and 1 subspaces with two triplets on j and k rungs are obtained as,

$$(j,k)_0 = \frac{2}{\sqrt{3}}\left(\langle j,k\rangle_\mathrm{l} - \frac{1}{2}\langle j,k\rangle_\mathrm{r}\right) \quad (13)$$

and

$$(j,k)_1 = \frac{1}{\sqrt{2}}\Big(\{j,k\}_\mathrm{u}[j,k]_\mathrm{l} + [j,k]_\mathrm{u}\{j,k\}_\mathrm{l}\Big), \quad (14)$$

respectively. To simplify expressions of the perturbation theory, we use the Fourier transformation of one- and two-triplet excited state given by,

$$|1 : q_x\rangle = \frac{1}{\sqrt{N}}\sum_j e^{iq_x j}\{j,j\}_\mathrm{i}\prod_{m\neq j}[m,m]_\mathrm{i} \quad (15)$$

and

$$|2 : S,(q_x,L)\rangle = \frac{1}{\sqrt{N}}\sum_j e^{iq_x j}(j,j+L)_S\prod_{m\neq j,j+L}[m,m]_\mathrm{i}, \quad (16)$$

respectively, where $S = 0, 1$ and $L = 1, 2, \cdots, (N-1)/2$.

III. RESULTS

In this section, we present numerical results of magnetic excitations in the frustrated two-leg spin-ladder system, and discuss the frustration effects by J_2. Figure 2 shows the DSCF in a 32-rung frustrated ladder for three phases: the incommensurate CD phase $(J_p/J_1 = 0.2,\ J_2/J_1 = 0.6)$, the incommensurate RS phase $(J_p/J_1 = 1.0,\ J_2/J_1 = 0.6)$, and the commensurate RS phase $(J_p/J_1 = 1.0,\ J_2/J_1 = 0.1)$.

First, we can find an obvious difference between the CD phase and the RS phase in the DSCF. That is, the spectrum of the bonding mode in the CD phase (Fig. 2(a)) exhibits almost the same shape with the same intensity as that of the anti-bonding mode (Fig. 2(b)), while those are completely different in the RS phase (Fig. 2(c)-(f)). The difference between the CD phase and the RS phase is understood as the difference of the elementary excitations in both phases. In the CD phase, the key elementary excitation is spinon, which strongly reflects one dimensionality. Thus, there is few difference between the bonding and anti-bonding modes in the CD phase. On the other hand, the major elementary excitation in the RS phase is triplon, which is the excitations from singlet to triplet in rung bonds. The parity of the triplet state is different from that of the singlet state. Therefore, the excitation exchanging the rung parity, that is anti-bonding mode $(q_y = \pi)$, is dominant in the RS phase. With finite J_1, the ground state is mixed with singlet states in subspaces of even-number triplets, which has the same parity as the even-number triplet excitations. Hence, the excitation of the bonding mode $(q_y = 0)$ appears with finite J_1. We stress that the difference in the spectral weights plays a key role in identifying the ground-state phase by using inelastic-neutron-scattering data for the frustrated two-leg spin-ladder compound, $\mathrm{BiCu_2PO_6}$.

In the following subsections, we clarify the origin of the spectra for each phase with three excitations: spinon, triplon, and bound state of two triplons. The frustration effects are also clarified.

A. Multi-spinon excitations

The phase transition between the CD and RS phases in the frustrated ladder system belongs to the same universality class as the Ising transition[6]. The crucial point of the transition is a change of the ground-state degeneracy and the symmetry. Although the ground state has no degeneracy in the RS phase, there are twofold-degenerate ground states in the CD phase. At zero temperature, the translation symmetry is spontaneously broken in the CD phase. The CD phase continuously changes to the dimer phase in the J_1-J_2 model in the zero-coupling limit in the rung direction. Therefore, the key excitation should be spinon and is almost identical to the elementary excitation in the frustrated chain system.

In Figs. 2 (a) and (b), we can see the lower edge of



FIG. 2. (Color online) The dynamical spin correlation functions (DSCF) of a 32-rung frustrated ladder for three phases: (a), (b) the incommensurate CD phase ($J_p/J_1 = 0.2,\ J_2/J_1 = 0.6$), (c), (d) the incommensurate RS phase ($J_p/J_1 = 1.0,\ J_2/J_1 = 0.6$), and (e), (f) the commensurate RS phase ($J_p/J_1 = 1.0,\ J_2/J_1 = 0.1$). (a), (c), and (e) [(b), (d), and (f)] represent the excitations with $q_y = 0$ [$q_y = \pi$][25]. We set the DMRG truncation number $m = 150$ and the broadening factor $\gamma = 0.1$.

the two-spinon continuum, which corresponds to the des Cloizeaux-Pearson mode in the unfrustrated limit $J_2 \to 0$ in the chain system[11,26]. The excitation has a minimum energy at an incommensurate wavenumber q_x^* near the classical solution $q_{x(\mathrm{cl})}^* = \cos^{-1}(-J_1/4J_2) \cong q_x^*$. The dispersion relation is almost symmetric with respect to $q_x = \pi/2$, although the intensity is weaker for $q_x < \pi/2$ than $q_x > \pi/2$. This feature originates from the doubled unit cell due to the dimerization. To explain this briefly, we consider the one-dimensional version of the DSCF for the frustrated chain system at the Majumdar-Ghosh point $J_2/J_1 = 0.5$ with the periodic boundary condition. The ground states doubly degenerate as follows:

$$|\mathrm{MG}\rangle_l = \prod_{j=0}^{N/2-1} [2j, 2j+1], \tag{17}$$

$$|\mathrm{MG}\rangle_r = \prod_{j=0}^{N/2-1} [2j+1, 2j+2]. \tag{18}$$

We assume that the system size N is even. The ground-state energy of the Majumdar-Ghosh Hamiltonian is easily obtained as $\epsilon_{\mathrm{MG}} = -\frac{3}{8}NJ_1$. The one-dimensional version of the DSCF for the left MG state $|\mathrm{MG}\rangle_l$ is given by,

$$\chi(q,\omega)^{(\mathrm{1D})} = -\frac{1-\cos(q)}{2\pi}\Im\langle q|\frac{1}{\omega - (\mathcal{H}_{\mathrm{MG}} - \epsilon_{\mathrm{MG}}) + i0}|q\rangle, \tag{19}$$

with,

$$|q\rangle = \frac{1}{\sqrt{N}}\sum_{j=0}^{N/2-1} e^{2ijq}\{2j, 2j+1\}\prod_{m\neq j}[2m, 2m+1]. \tag{20}$$

Thus, we obtain the relation between $\chi(q,\omega)^{(\mathrm{1D})}$ and $\chi(\pi-q,\omega)^{(\mathrm{1D})}$ as,

$$\chi(\pi-q,\omega)^{(\mathrm{1D})} = \frac{1+\cos(q)}{1-\cos(q)}\chi(q,\omega)^{(\mathrm{1D})}, \tag{21}$$

where $q \neq 0$. For $q = [\pi/2, \pi]$, we can easily confirm the relation about the spectral weight as, $\chi(\pi-q,\omega)^{(\mathrm{1D})} \leq \chi(q,\omega)^{(\mathrm{1D})}$, with the same poles. This is due to the fact that the doubled periodicity of the dimer state, which occurs spontaneously, folds the Brillouin zone in half. In other words, the original dispersion relation without dimerization and its inverted one with respect to $q = [\pi/2, \pi]$ are overlapped each other in the Brillouin zone. Figures 2 (a) and (b) have such a situation.

B. Triplon excitation

We discuss the anti-bonding mode ($q_y = \pi$) in the RS phase (Fig. 2 (d) and (f)) in terms of *triplon*. The triplon excitation is defined as the rung triplet excitations in this study. In the perturbation theory from the strong coupling limit in the rung direction, the dispersion relation of the triplon within the second order is obtained as

$$\epsilon_{\rm T}(q_x) = J_p \left[1 + \alpha_1 \cos(q_x) + \alpha_2 \cos(2q_x) + \frac{3}{4}\left(\alpha_1^2 + \alpha_2^2\right) + O(\alpha^3) \right], \tag{22}$$

where $\alpha_1 \equiv J_1/J_p$ and $\alpha_2 \equiv J_2/J_p$. The wavenumber at the minimum energy of the triplon excitation q_x^* corresponds to the incommensurability determined by the classical solution $q_{x({\rm cl})}^*$.

In Fig. 2 (d), the wavenumber of the lowest excitation $q_x^* = (2/3 \pm 1/33)\pi$ is obtained for the 32-rung ladder system. The wavenumber is slightly shifted from the classical one $q_{x({\rm cl})}^* \cong 0.637\pi$. It is known that the quantum correction to the wavenumber can be obtained by a mean-field analysis with rung-bond operators. However, the analysis requires proper renormalization to obtain a precise energy, which is, however, demanding. Thus, the numerical calculation is indispensable to obtain the precise wavenumber with the minimum energy in the dispersion relation of the triplon, especially, for large α_1 and α_2 region.

In Fig. 2 (d), we find that the intensity is concentrated at the wavenumber $q \sim 2\pi/3$ with the minimum energy. In Fig. 2 (f), both the maximum intensity and the minimum energy shift from $q \sim 2\pi/3$ to $q = \pi$ simultaneously. Thus, we conclude that the frustration affects only the wavenumber and the triplon picture is still stable.

C. Bound state of two triplons

In this subsection, we discuss the bonding mode ($q_y = 0$) in the RS phase (Figs. 2 (c) and (e)). First, we show that the bonding mode disappears in the strong coupling limit $J_p/J_1 \to \infty$ in the rung direction without frustration $J_2 = 0$. The ground state in this limit is given by the direct product of the rung singlets, $|0\rangle^{(0)} = \prod_j [j,j]_{\rm i}$. In addition, the excited state at $q_y = 0$ in Eq. (5), $S^z(q_x, q_y = 0)|0\rangle^{(0)}$, is explicitly rewritten as

$$S^z(q_x, 0)|0\rangle^{(0)} = \left(\frac{1}{\sqrt{N}} \sum_j e^{iq_x j} S^z_{j,q_y=0} \right) \left([j,j]_{\rm i} \prod_{m \neq j} [m,m]_{\rm i} \right) = 0, \tag{23}$$

because $S^z_{j,q_y=0}[j,j]_{\rm i} = 0$. Therefore, the intensity of the bonding mode should vanish in the strong coupling limit in the rung direction.

Next, we consider the perturbation effect of $\alpha_1 = J_1/J_p \ll 1$ on $S = 1$ excitation in the bonding mode. The ground state within the first order of the perturbation with respect to α_1 is obtained as,

$$|0\rangle^{(1)} = -\frac{\sqrt{3}}{4}\alpha_1 \sqrt{N} |2:0,(0,1)\rangle. \tag{24}$$

The excited state from this state exists: $S^z(q_x, 0)|0\rangle^{(1)} \neq 0$. For example, the excited state at $q_x = \pi$ is obtained as,

$$S^z(\pi, 0)|0\rangle^{(1)} = -\frac{\alpha_1}{\sqrt{2}} |2:1,(\pi,1)\rangle. \tag{25}$$

This is the origin of the bonding mode, so-called *bound triplon*. Since the bound triplon appears proportional to the first order of α_1, the DSCF of the bound triplon starts from the second order. Figure 3 shows the α_1 dependence of the bound triplon at $q_x = \pi$ obtained by the dynamical DMRG calculation for a 32-rung unfrustrated ladder with the truncation number $m \geq 300$. In the inset of Fig. 3 (b), we can see that the peak energy is proportional to α_1^2.

We can obtain the excitation energy of the $S = 1$ state of two triplons at $q_x = \pi$ by the perturbation theory. In addition to the bound triplon whose energy is $\epsilon_{\rm BT}/J_p = 2 - \frac{1}{2}\alpha_1 + \frac{9}{8}\alpha_1^2 + O(\alpha_1^3)$ (Fig. 3 (b)), there is a continuum of two triplons starting from the energy, $\epsilon_l/J_p = 2 + \frac{9}{8}\alpha_1^2 + O(\alpha_1^3)$. In fact, we can see the continuum above the bound triplon (the inset of Fig. 3 (a)). We note that the bound triplon is stabilized because of the dimerization energy in the leg, $-\frac{3}{4}J_1 + \frac{1}{4}J_1 = -\frac{1}{2}J_1$.

The frustration effect is also investigated with the perturbation theory. Figure 4 shows the α_2 dependence of DSCF at $q_x = \pi$ with $\alpha_1 = 0.5$. We can see that the energy of the bound triplon shifts to the continuum of two triplons as α_2 increases. At the same time, the weight of the bound triplon moves into the continuum. Within the second order of α_1 and in the first order of α_2, i.e., $\alpha_2/\alpha_1 \ll 1$ is assumed, $\epsilon'_{\rm BT}/J_p = 2 - \frac{1}{2}\alpha_1 + \frac{9}{8}\alpha_1^2 + \alpha_2 + O(\alpha_1^3, \alpha_2^2, \alpha_1\alpha_2)$ at $q_x = \pi$ by the perturbation analysis. In the inset of Fig. 4, we can make sure that the energy of the bound triplon follows the equation. We expect the collective mode to be weakened and to be buried in the continuum with large J_2/J_1 in the strong coupling limit in the rung direction. Therefore, the bound triplon is unstable with large frustration, while the triplon is still stable with a slight shift of the wavenumber.



FIG. 3. (Color online) (a) α_1 dependence of the DSCF in a 32-rung unfrustrated ladder, namely $J_2 = 0$, at $(q_x, q_y) = (32\pi/33, 0)$ in the commensurate RS phase. The inset of (a) shows the DSCF for $\alpha_1 = 1/2$ (red solid line) and Lorentzian fitting (green dotted line). (b) The energy of the peak versus α_1 where the energy of the bound triplon with second order perturbation theory, $\epsilon_{\mathrm{BT}}^{(2)}/J_p = 2 - \alpha_1/2 + 9\alpha_1^2/8$. The inset of (b) shows the DSCF at the peak versus α_1. Blue line shows a fitting function proportional to α_1^2 with the coefficient, 0.265 ± 0.002. We set the DMRG truncation number $m = 300$ and the broadening factor $\gamma = 0.05$ to distinguish the high-energy continuum from the main intensity of single Lorentzian.



FIG. 4. (Color online) α_2 dependence of the DSCF in a 32-rung frustrated ladder at $(q_x, q_y) = (32\pi/33, 0)$ at $\alpha_1 = 0.5$. The inset shows the energy of the peak, corresponding to the energy of the bound triplon, as a function of α_2. Here, $\epsilon'^{(1)}_{\mathrm{BT}} = \epsilon_{\mathrm{BT}}(\alpha_1) + \alpha_2 J_p$ (green line). We set the DMRG truncation number $m = 300$ and the broadening factor $\gamma = 0.05$

IV. SUMMARY AND DISCUSSION

The frustrated two-leg spin-ladder system was investigated as a quantum spin system bridging two different systems: the frustrated chain system and the unfrustrated ladder system. We examined the dynamical spin correlation function at zero temperature to understand the elementary excitations and the frustration effects, by using dynamical density-matrix renormalization-group method. We showed the excitation spectra for three phases: the incommensurate columnar-dimer phase, the incommensurate rung-singlet phase and the commensurate rung-singlet phase.

First, we made clear that the q_y dependence of the spectrum is different between the columnar-dimer phase and the rung-singlet phase. This is because the elementary excitations are different between the two phases. A key excitation in the columnar-dimer phase is spinon, which is the same as the frustrated chain system. In contrast, a major excitation is triplon in the rung-singlet phase. The difference was explained by using both spinon picture for the columnar-dimer phase and triplon picture for the rung-singlet phase. With the help of the perturbation theory from the strong coupling limit in the rung direction, we clarified the instability of the bound triplon with the frustration, although the triplon is still stable.

This is different from a frustration effect on the triplon excitations in the ladder system[16], where the frustration is introduced with a diagonal antiferromagnetic interaction (J_d) in the nearest-neighbor-bond square. This system corresponds to a frustrated chain with alternating the nearest-neighbor interaction[27–30]. The RS phase appears without J_d, and the dimer phase appears at $J_d = J_1$ with large J_1 in the frustrated ladder system. Therefore, from a viewpoint on the ground-state phase transition, the in-chain frustration (J_2) effect is similar to the diagonal frustration (J_d) effect for the ladder system. However, the frustration effects on the excitations are different, because the triplon excitation is fragile with the diagonal frustration, but not with the in-chain frustration as shown in the present study. Thus, the frustration effects on the excitations are important for clarifying the influence of the in-chain frustration in the ladder system.

Experimentally, this model will be useful for understanding the low-energy physics in a newly synthesized compound $\mathrm{BiCu_2PO_6}$. The magnetic structure of $\mathrm{BiCu_2PO_6}$ is characterized by $\frac{1}{2}$ spins on Cu sites arranged in zigzag chains in the b axis[19–21]. The zigzag chains are stacked up in the c axis. The exchange interactions J_1 and J_2 are defined as the nearest-neighbor and next-nearest-neighbor interactions in the zigzag chain. In addition, there are comparable inter-chain interactions in $\mathrm{BiCu_2PO_6}$. Since the distance between chains is alternate, we define the rung exchange interactions J_p and J'_p as the nearest-neighbor interactions on the shorter and longer bonds, respectively. The compound was reported to be similar to a unfrustrated ladder system at first. That is, J_2 and J_4 are negligible as compared with J_1. However, the following numerical studies have shown that the susceptibility can be understood well as that in a frustrated two-leg spin-ladder system, where the frustration is introduced as a next-nearest-neighbor antiferromagnetic interaction. Namely, the frustration, J_2, should not be ignored. Besides, the band-structure calculation presented by Tsirlin *et al.*[21] has claimed that J'_p interaction is comparable to J_1 and J_2, although J_p is rather negligible. By comparing our results and inelastic neutron scattering experiments[25], the magnitude of magnetic exchange interactions and the ground state of such a compound will be determined in the near future.

ACKNOWLEDGMENTS

We thank M. Fujita and R. Kajimoto for fruitful discussions. Special thanks also go to T. Adachi, K. Ishii, K. Ikeuchi, K. Tsutsui, and T. Sakai who gave me invaluable comments. This work was partly supported by Grant-in-Aid for Scientific Research from MEXT (Grant No.24540387, No.24360036, No.23340093, and No.22340097) and by the inter-university cooperative research program of IMR, Tohoku University. We also acknowledge the support from the Strategic Programs for Innovative Research (SPIRE), the Computational Materials Science Initiative (CMSI) and the Global COE Program "The Next Generation of Physics, Spun from Universality and Emergence", Kyoto University. Numerical computation in this work was carried out on the supercomputers at JAEA, YITP in Kyoto University and ISSP in University of Tokyo.

Appendix: Perturbation theory with RVB picture

We describe the perturbation-theory analysis in the rung-singlet phase for the two-leg spin-ladder system. Because of the symmetries of the Hamiltonian, i.e., translation in the leg direction and reflection of two legs, RVB picture is useful for analyzing low-energy states with perturbation theory. Some simple relations helps us to calculate high-order states and the energies with perturbation theory. In this paper, we concentrate on the ground state, one-triplet excitations, and two-triplet excitations.

First, we show some equations regarding to the rung-singlet configuration. In the rung-singlet phase, the ground state is described as a linear combination of some leg-singlet-pair states. In practice, the perturbed Hamiltonian acting on the rung-singlet configuration generates a leg-singlet-pair state:

$$h_j^{(R)}\langle j, j+R\rangle_{\mathrm{r}} = \frac{1}{2}\langle j, j+R\rangle_{\mathrm{r}} - \langle j, j+R\rangle_{\mathrm{l}}, \tag{A.1}$$

where $R = 1, 2$. In addition, the perturbed Hamiltonian can generate a long-range leg-singlet pair and crossing leg-singlet pairs as follows,

$$h_j^{(R)}[j,j]_{\mathrm{i}}\langle j+R, k\rangle_{\mathrm{l}} = \frac{1}{2}[j+R, j+R]_{\mathrm{i}}\langle j,k\rangle_{\mathrm{l}} - \frac{1}{2}[k,k]_{\mathrm{i}}\langle j, j+R\rangle_{\mathrm{l}}, \tag{A.2}$$

$$h_j^{(R)}\langle i,j\rangle_{\mathrm{l}}\langle j+R, k\rangle_{\mathrm{l}} = \frac{1}{2}\langle i, j+R\rangle_{\mathrm{l}}\langle j,k\rangle_{\mathrm{l}} - \frac{1}{2}\langle i,k\rangle_{\mathrm{l}}\langle j, j+R\rangle_{\mathrm{l}}. \tag{A.3}$$

Within low-order perturbations, the corrections of the ground state and the excited states can be easily obtained with these equations.

For the one-triplet excited state, the perturbed Hamiltonian can only move the triplet to another rung without generating leg-singlet pairs:

$$h_j^{(R)}\{j,j\}_{\mathrm{i}}[j+R, j+R]_{\mathrm{i}} = \frac{1}{2}[j,j]_{\mathrm{i}}\{j+R, j+R\}_{\mathrm{i}}, \tag{A.4}$$

$$h_j^{(R)}\{j,j\}_{\mathrm{i}}\langle j+R, k\rangle_{\mathrm{l}} = \frac{1}{2}\{j+R, j+R\}_{\mathrm{i}}\langle j,k\rangle_{\mathrm{l}} - \frac{1}{2}\{k,k\}_{\mathrm{i}}\langle j, j+R\rangle_{\mathrm{l}}. \tag{A.5}$$

These equations means that the frustration never make a rung triplet unstable. In particular, Eq. (A.4) determines the dispersion relation of the triplon in Eq. (22).

On the other hand, the perturbed Hamiltonian acting on a leg-triplet state generates a longer-range leg-triplet state as follows,

$$h_j^{(R)}[j,j]_{\mathrm{i}}(j+R,k)_1 = \frac{1}{2}[j+R,j+R]_{\mathrm{i}}(j,k)_1, \tag{A.6}$$

$$h_j^{(R)}\langle i,j\rangle_{\mathrm{l}}(j+R,k)_1 = \frac{1}{2}\langle i,j+R\rangle_{\mathrm{l}}(j,k)_1 - \frac{1}{2}\langle j,j+R\rangle_{\mathrm{l}}(i,k)_1. \tag{A.7}$$

This means that the bound state, i.e., a nearest-neighbor leg-triplet state, is no longer stable with any perturbation. With only nearest-neighbor leg interaction $h_j^{(1)}$, off-diagonal parts of the perturbed Hamiltonian for the momentum distribution of leg-triplet state are canceled out at $q = \pi$. This is the reason why the bound triplon is stable around $q = \pi$. The next-nearest-neighbor leg interaction $h_j^{(2)}$ mixes the nearest-neighbor leg-triplet state with longer-range leg-triplet states. Thus, the bound triplon is smeared out with large frustration.

---

[*] sugimoto.takanori@jaea.go.jp

[1] As a review, S. Maekawa, T. Tohyama, S. E. Barnes, S. Ishihara, W. Koshibae, and G. Khaliullin, *Physics of Transition Metal Oxides*, vol. 144 of *Springer Series in Solid-State Sciences*, Springer (2004).

[2] C. Kim, A. Y. Matsuura, Z.-X. Shen, N. Motoyama, H. Eisaki, S. Uchida, T. Tohyama, and S. Maekawa, Phys. Rev. Lett. **77**, 4054 (1996).

[3] B. Lake, A. M. Tsvelik, S. Notbohm, D. A. Tennant, T. G. Perring, M. Reehuis, C. Sekar, G. Krabbes, and B. Büchner, Nature Physics **6**, 50 (2009).

[4] As a review, L. Balents, Nature **464**, 199 (2010).

[5] K. P. Schmidt and G. S. Uhrig, Phys. Rev. Lett. **90**, 227204 (2003).

[6] A. Lavarélo, G. Roux, and N. Laflorencie, Phys. Rev. B, **84**, 144407 (2011).

[7] I. Affleck, D. Gepner, H. J. Schulz, and T. Ziman, J. Phys. A **22**, 511 (1989).

[8] K. Okamoto and K. Nomura, Phys. Lett. A **169**, 433 (1992).

[9] S. Eggert, Phys. Rev. B **54**, R9612 (1996).

[10] G. Müller, H. Thomas, H. Beck, and J. C. Bonner, Phys. Rev. B **24**, 1429 (1981).

[11] H. Yokoyama and Y. Saiga, J. Phys. Soc. Jpn. **66**, 3617 (1997).

[12] C. K. Majumdar and D. K. Ghosh, J. Math. Phys. **10**, 1399 (1969).

[13] S. R. White and I. Affleck, Phys. Rev. B **54**, 9862 (1996).

[14] T. Barnes, E. Dagotto, J. Riera, and E. S. Swanson, Phys. Rev. B **47**, 3196 (1993).

[15] S. Gopalan, T. M. Rice, and M. Sigrist, Phys. Rev. B **49**, 8901 (1994).

[16] O. P. Sushkov and V. N. Kotov, Phys. Rev. Lett. **81**, 1941 (1998); V. N. Kotov, O. P. Sushkov, and R. Eder, Phys. Rev. B **59**, 6266 (1999).

[17] N. Haga and S.-I. Suga, Phys. Rev. B **66**, 132415 (2002).

[18] F. Abraham, M. Ketatni, G. Mairesse, and B. Mernari, Eur. J. Solid State Inorg. Chem. **31**, 313 (1994).

[19] B. Koteswararao, S. Salunke, A. V. Mahajan, I. Dasgupta, and J. Bobroff, Phys. Rev. B **76**, 052402 (2007).

[20] O. Mentré, E. Janod, P. Rabu, M. Hennion, F. Leclercq-Hugeux, J. Kang, C. Lee, M.-H. Whangbo, and S. Petit, Phys. Rev. B **80**, 180413(R) (2009).

[21] A. A. Tsirlin, I. Rousochatzakis, D. Kasinathan, O. Janson, R. Nath, F. Weickert, C. Geibel, A. M. Läuchli, and H. Rosner, Phys. Rev. B **82**, 144426 (2010).

[22] S. Notbohm, P. Ribeiro, B. Lake, D. A. Tennant, K. P. Schmidt, G. S. Uhrig, C. Hess, R. Klingeler, G. Behr, B. Büchner, M. Reehuis, R. I. Bewley, C. D. Frost, P. Manuel, and R. S. Eccleston, Phys. Rev. Lett. **98**, 027403 (2007).

[23] S. R. White, Phys. Rev. Lett. **69**, 2863 (1992); Phys. Rev. B **48**, 10345 (1993).

[24] E. Jeckelmann, Phys. Rev. B **66**, 045114 (2002).

[25] We note that the Brillouin zone in the leg direction for $\mathrm{BiCu_2PO_6}$ is half of our theoretical one because the unit cell in the compound includes two rungs. Thus, the spectrum in Fig. 2 will be folded in half to be compared with experimental one.

[26] T. Sugimoto, S. Sota, and T. Tohyama, J. Phys. Soc. Jpn. **81**, 034706 (2012).

[27] S. Watanabe and H. Yokoyama, J. Phys. Soc. Jpn. **68**, 2073 (1999).

[28] N. Haga and S.-I. Suga, Phys. Rev. B **65**, 014414 (2001).

[29] T. Vekua and A. Honecker, Phys. Rev. B **73**, 214427 (2006).

[30] The interactions in the frustrated ladder system, J_1, J_p, and J_d, correspond to the interactions in the frustrated chain system with alternation, J_2, J_1, and J_1', respectively.